



every child by two!

**Children Need Vaccinations  
at the Following Ages:**

- Birth
- 2 Months
- 4 Months
- 6 Months
- 12 - 15 Months
- 4 - 6 Years
- 11 - 12 Years

Additionally, influenza (flu) vaccine is recommended every year for children 6 months through 18 years.

Vaccines recommended by the Advisory Committee on Immunization Practices protect children against diphtheria, tetanus (lock jaw), polio, measles, pertussis (whooping cough), mumps, rubella (german measles), haemophilus influenzae type b, hepatitis A & B, varicella (chickenpox), invasive pneumococcal disease, invasive meningococcal disease, rotavirus and influenza (flu).

Be sure to complete any series of immunizations once started. One or two doses of a series is not adequate protection. Older children may need additional vaccines or booster shots. Call your doctor or local health clinic for more information.

The following chart lists the names of the vaccines recommended for your child and the diseases that they prevent.

Disease Name	Vaccine Name
Measles	Measles Vaccine (contained in MMR vaccine)
Mumps	Mumps Vaccine (contained in MMR vaccine)
Rubella (German Measles)	Rubella Vaccine (contained in MMR vaccine)
Diphtheria	Diphtheria Toxoid (contained in DTP, DTaP, DT, and TD vaccines)
Tetanus (Lock Jaw)	Tetanus Toxoid (contained in DTP, DTaP, DT, or TD vaccines)
Pertussis (Whooping Cough)	Pertussis Vaccine (contained in DTP and DTaP vaccines)
Hepatitis B	Hepatitis B Vaccine (also known as Hep B vaccine)
Hib	Haemophilus Influenzae type b (Hib) Vaccine
Influenza (Flu)	Influenza Vaccine
Polio	Inactivated Polio Vaccine (also known as IPV vaccine)
Varicella (Chickenpox)	Varicella Vaccine
Pneumococcal Disease	Pneumococcal Conjugate Vaccine (also known as the PCV vaccine)
Hepatitis A	Hepatitis A Vaccine (also known as Hep A vaccine)
Meningococcal disease/Meningitis	Meningococcal Conjugate Vaccine
Rotavirus	Rotavirus Vaccine (RV)